

# Simple Procedure for the Designation of Haar Wavelet Matrices for Differential Equations

Phang Chang, Phang Piau

**Abstract** — Wavelet transforms or wavelet analysis is a recently developed mathematical tool for many problems. Wavelets also can be applied in numerical analysis. In this paper, we apply Haar wavelet methods to solve ordinary differential equations with initial condition known. To avoid the tedious calculations and to promote the study of wavelets to beginners, we proposed a simple way to perform the calculations for the matrix representation. Three numerical examples are shown which including first, second, higher order differential equations with constant and variable coefficients. The results show that the proposed way are quite reasonable when compare to exact solution.

**Index Terms** — Haar wavelet methods, matrix representation, ordinary differential equations, computer algebra system

## I. INTRODUCTION

Wavelet transform or wavelet analysis is a recently developed mathematical tool for signal analysis. To date, wavelets have been applied in numerous disciplines such as image compression, data compression, denoising data and many more [1]. In numerical analysis, wavelets also serve as a Galerkin basis to solve partial differential equations. Wavelet analysis involves tedious calculations. Practically, the calculation is done by using software with certain commands or special toolboxes. It may make the beginner feel intimidate [2]. Meanwhile, Haar function always has been choose for educational purpose, especially in many papers or books written on topic of introduction to wavelets [3-4]. Due to the powerful of wavelets in plenty fields, there are many works, such as in [5-6] in promoting the study of wavelet even in undergraduate level.

Meanwhile in numerical analysis, wavelet based algorithms have become an important tools because of the properties of localization. One of the popular families of wavelet is Haar wavelets. Due to its simplicity, Haar wavelets had become an effective tool for solving many problems, among that are Ordinary Differential Equations, ODEs and Partial Differential Equations, PDEs. In these works, the operator or matrix representation is expanded in a wavelet

Manuscript received December 12, 2007. This work was partially supported in part by the University Tun Hussein Onn Malaysia under Short Term Grant vot0356.

Phang Chang is with the Centre for Science Studies, University Tun Hussein Onn Malaysia, UTHM. 86400 Batu Pahat, Johor, Malaysia (phone: 607-4537973 ; fax: 607-4536051; e-mail: pchang@uthm.edu.my).

Phang Piau is with Department of Computational Science and Mathematics, Faculty of Computer Science and Information Technology, University Malaysia Sarawak, UNIMAS, 94300 Kota Samarahan, Sarawak, Malaysia. (e-mail: pphang@fit.unimas.my).

basis. Sometime, the works for writing up the operational matrices are quite tedious especially when ones intend to perform the calculation in high resolution. This will discourage the beginner to study how wavelet basis can be applied to solve differential equations, especially when in the works of encouraging the study of wavelets in undergraduate level which were done in [5-7].

In solving ordinary differential equations by using Haar wavelet related method, Chen and Hsiao [8-9] had derived an operational matrix of integration based on Haar wavelet. Lepik [10] had solved higher order as well as nonlinear ODEs by using Haar wavelet method. There are discussions by other researchers [11-13]. We are not going to compare with these distinguish scholars, but intend to come out with a simple procedure to solve ordinary differential equations which make use of the power of wavelets. With this, we hope that even for an undergraduate student can also perform the numerical calculation by using wavelets as a tool in order to solve ordinary differential equation problems without using any complicated algorithm but sufficient with the help of computer algebra system or Excel. With this, it is hoped that it will encourage the study of wavelet in undergraduate level.

## II. HAAR WAVELET

Haar wavelet is the simplest wavelet. Haar transform or Haar wavelet transform has been used as an earliest example for orthonormal wavelet transform with compact support. The Haar wavelet transform is the first known wavelet and was proposed in 1909 by Alfred Haar. The Haar function can be described as a step function  $\psi(x)$  and in Fig. 1 as follows :

$$\psi(x) = \begin{cases} 1 & 0 \leq x \leq 0.5 \\ -1 & 0.5 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (1)$$




Fig. 1 Haar function

This is also called mother wavelet. In order to perform wavelet transform, Haar wavelet uses translations and dilations of the function, i.e. the transform make use of following function

$$\psi(x) = \psi(2^j x - k) \quad (2)$$

Translation / shifting  $\psi(x) = \psi(x - k)$

Dilation / scaling  $\psi(x) = \psi(2^j x)$

where this is the basic works for wavelet expansion.

With the dilation and translation process as in Eq.(2), ones can easily obtain father wavelet, daughter wavelet, granddaughter wavelet and so on as in fig. 2.




Fig. 2 Haar Wavelet (up to 2 resolution levels)

In the matrix form, the Haar matrix for resolution up to 2 levels is given below :

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & -1 & -1 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

The pattern of the Haar wavelet undergo translation and dilation process and its matrix pattern are observed when the Haar wavelet been used in solving ordinary differential equations.

### III. HAAR WAVELET METHOD FOR ORDINARY DIFFERENTIAL EQUATIONS

For solving linear ordinary differential equation with  $n^{\text{th}}$  order, say

$$A_1 y^{(n)}(x) + A_2 y^{(n-1)}(x) + \dots + A_n y(x) = f(x),$$

where  $x \in [A, B]$  and initial conditions

$$y^{(n-1)}(A), y^{(n-2)}(A) \dots, y(A) \text{ are known.}$$

We follow the work done by Lepik [10]. Say we intend to do until  $j$  level of resolution, hence we let  $m = 2(2^j)$ . The interval  $[A, B]$  will be divided into  $m$  subintervals, hence  $\Delta x = \frac{B-A}{m}$  and the matrices are in the dimension of  $m \times m$ .

Here we suggest the step by step procedures for easy understanding. Mainly, there are 5 steps as shown in the procedure as follow.

Procedure :

Step 1: Let  $y^{(n)}(x) = \sum_{i=1}^m a_i h_i(x)$  where  $h$  is haar matrix and

$a_i$  is the wavelet coefficients.

Step 2: Obtain appropriate  $v$  order of  $y(x)$  by using

$$y^{(v)}(x) = \sum_{i=1}^m a_i P_{n-v,i}(x) + \sum_{\sigma=0}^{n-v-1} \frac{1}{\sigma!} (x-A)^\sigma y_0^{(v+\sigma)}$$

Step 3: Replace  $y^{(n)}(x)$  and all the value of  $y^{(v)}(x)$  into the problem.

Step 4: Calculate the wavelet coefficients,  $a_i$ .

Step 5: Obtain the numerical solution for  $y(x)$ .

Step 2 is the key procedure where matrix  $P_{n-v,i}(x)$  will be counted. If ones intend to do the calculation until level  $j$  of resolution, ones will obtain the matrix  $P_{n-v,i}(x)$  (let  $n-v = \alpha$ ) as in the pattern shown in fig. 3 where  $C = B - A$




Fig. 3: Designation of matrix  $P$ .

The pattern of design of the matrices  $P$  is similar to Haar wavelet. For calculation of  $P$  matrices, we focus on the elements need to be counted as shown in fig. 3. Here, we suggest the following algorithm for counting the elements which are required.

Algorithm :

$$\frac{1}{\alpha!} \frac{1}{(2m)^\alpha} \left[ (C \frac{m}{2^L} + 2l - 1)^\alpha - 2(C(2l - 1))^\alpha \right]$$

$$\alpha = n - v$$

$$L = 0, 1, 2, \dots, j \text{ (Level of Haar wavelet)}$$

$$l = 1, 2, 3, \dots, \frac{m}{2(2^L)}$$

and where  $C = B - A$

In the matrix  $P$  shown in the fig 3, we factored out the common factor  $\frac{1}{\alpha!} \frac{C^\alpha}{(2m)^\alpha}$  for all the elements in the matrix

which we calculated using the algorithm above, for the reason of simplicity.

IV. NUMERICAL EXAMPLES

Three examples are shown. All the calculation is done by using 3 levels of Haar wavelet.

Example 1:

Solve the equation  $y''(x) + y(x) = \sin x + x \cos x$ ,

$$x \in [0,1] \text{ with } y(0) = 1, y'(0) = 1.$$

Exact solution :

$$y(x) = \cos x + \frac{5}{4} \sin x + \frac{1}{4} (x^2 \sin x - x \cos x)$$

By using

$$\text{Step 1 : } y''(x) = \sum_{i=1}^m a_i h_i(x)$$

$$\begin{aligned} \text{Step 2 : } y(x) &= \sum_{i=1}^m a_i P_{2,i}(x) + \sum_{\sigma=0}^{2-0-1} \frac{1}{\sigma!} (x-0)^\sigma y_0^{(\sigma)} \\ &= \sum_{i=1}^m a_i P_{2,i}(x) + 1 + x \end{aligned}$$

Step 3 : Hence, we get

$$y''(x) + y(x) = \sin x + x \cos x$$

$$\sum_{i=1}^m a_i h_i(x) + \sum_{i=1}^m a_i P_{2,i}(x) + 1 + x = \sin x + x \cos x$$

$$\sum_{i=1}^m a_i [h_i(x) + P_{2,i}(x)] = \sin x + x \cos x - 1 - x$$

The matrix  $P$  is shown below.

$$\frac{1}{2!32^2} \begin{pmatrix} 1 & 3^2 & 5^2 & 7^2 & 9^2 & 11^2 & 13^2 & 15^2 & 17^2 & 19^2 & 21^2 & 23^2 & 25^2 & 27^2 & 29^2 & 31^2 \\ 1 & 3^2 & 5^2 & 7^2 & 9^2 & 11^2 & 13^2 & 15^2 & 287 & 343 & 391 & 431 & 463 & 487 & 503 & 511 \\ 1 & 3^2 & 5^2 & 7^2 & 79 & 103 & 119 & 127 & & & & & & & & \\ & & & & & & & & 1 & 3^2 & 5^2 & 7^2 & 79 & 103 & 119 & 127 \\ 1 & 3^2 & 23 & 31 & & & & & & & & & & & & \\ & & & & & 1 & 3^2 & 23 & 31 & & & & & & & \\ & & & & & & & & 1 & 3^2 & 23 & 31 & & & & \\ \frac{1}{2!32^2} & 1 & 7 & & & & & & & & & & 1 & 3^2 & 23 & 31 \\ & & & 1 & 7 & & & & & & & & & & & \\ & & & & & 1 & 7 & & & & & & & & & \\ & & & & & & & 1 & 7 & & & & & & & \\ & & & & & & & & & 1 & 7 & & & & & \\ & & & & & & & & & & & 1 & 7 & & & \\ & & & & & & & & & & & & & 1 & 7 & \\ & & & & & & & & & & & & & & & 1 & 7 \end{pmatrix}$$

Step 4: Solving the system of linear equation. We obtain wavelet coefficients,  $a_i$ .

Step 5: Obtain the numerical solution for  $y(x)$  as in Table 1.

Table 1 : Numerical solution for Example 1

$x (/32)$	Solution	Exact solution	Absolute error
1	1.030777	1.030767	0.000010
3	1.089527	1.089496	0.000031
5	1.144880	1.144700	0.000180
7	1.196844	1.196643	0.000201
9	1.247001	1.245594	0.001407
11	1.293240	1.291819	0.001421
13	1.337136	1.335577	0.001559
15	1.378678	1.377118	0.001560
17	1.427640	1.416676	0.010964
19	1.465381	1.454467	0.010914

21	1.501598	1.490681	0.010917
23	1.536265	1.525485	0.010780
25	1.570336	1.559012	0.011324
27	1.602462	1.591364	0.011098
29	1.633465	1.622605	0.010860
31	1.663308	1.652763	0.010545

Example 2 shows the solution of higher order differential equation by using Haar wavelet method. The result shows that the method is applicable for higher order DE, even for variable coefficient.

Example 2:

Solve the equation  $y^{(4)}(x) + xy(x) = 16 \sin 2x + x \sin 2x$ ,

$$x \in [0,1] \text{ with } y(0) = 0, y'(0) = 2, y''(0) = 0, y'''(0) = -8$$

Exact solution :  $y(x) = \sin 2x$

By carry out Step 1 to 3, we obtain

$$\sum_{i=1}^m a_i [h_i(x) + x P_{4,i}(x)] + 2x^2 - \frac{4}{3} x^4 = 16 \sin 2x + x \sin 2x$$

By carry out Step 4 to 5, we obtained the results as in Table 2.

Table 2 : Numerical solution for Example 2

$x (/32)$	Solution	Exact solution	Absolute error
1	0.062459	0.062459	0.000000
3	0.186405	0.186403	0.000002
5	0.307463	0.307439	0.000024
7	0.423743	0.423676	0.000067
9	0.533859	0.533303	0.000556
11	0.635637	0.634607	0.001030
13	0.727687	0.726009	0.001678
15	0.808581	0.806081	0.002500
17	0.887019	0.873575	0.013444
19	0.946546	0.927437	0.019109
21	0.992728	0.966827	0.025901
23	1.024951	0.991129	0.033822
25	1.042978	0.999966	0.043012
27	1.046518	0.993198	0.053320
29	1.035738	0.970932	0.064806
31	1.010991	0.933514	0.077477

Example 3 involves ODE with exponential coefficients. The results is compare with the results obtained by series solution. Due to the behavior of exponential, we use a relatively small step size if compared to previous two examples, which is until 4 level of resolution.

Example 3:

Solve the equation  $y'(x) + e^x y(x) = x^2$ ,

$$x \in [0,1] \text{ with } y(0) = 4.$$

Series solution :  $y(x) = 4 - 4x + x^3 + \frac{1}{12} x^4$  (up to first four nonzero terms only)

By carry out Step 1 to 3, we obtain

$$y'(x) + e^x y(x) = \sum_{i=1}^m a_i [h_i(x) + e^x P_{1,i}(x)] = x^2 - 4e^x$$

By carry out Step 4 to 5, we obtained the results as in Table 3

Table 3 : Numerical solution for Example 3

$x$ (/64)	Solution	Series solution	Difference
1	3.936517	3.937504	0.000987
3	3.807570	3.812603	0.005034
5	3.674638	3.687980	0.013342
7	3.537674	3.563820	0.026146

## V. CONCLUSION

The designation of matrices  $P$  is shown. The algorithm and procedure have been applied to use Haar wavelet method in solving ODEs. The result is comparable to the exact solution.

## REFERENCES

- [1] A.W. Galli, G.T. Heydt, and P.F. Ribeiro, "Exploring the power of wavelet analysis." IEEE Computer Application in Power, Oct 1996, pp.37 – 41.
- [2] M.N.O. Sadiku, C.M. Akujuobi, and R.C. Garcia, "An introduction to wavelets in electromagnetics." Microwave Magazine, IEEE, vol. 6(2), June 2005, pp. 63 – 72.
- [3] G.P. Nason, "A little introduction to wavelets." Applied Statistical Pattern Recognition, IEEE Colloquium on 20 April 1999, pp.1 – 6.
- [4] E. Aboufadel and S. Schlicker, *Discovering Wavelets*, New York : John Wiley & sons, Inc, 1999, pp. 12-18.
- [5] M.C. Potter, J.L. Goldberg, E.F. Aboufadel, *Advanced Engineering Mathematics*. 3<sup>rd</sup> Ed. New York : Oxford University Press. pp. 670-699.
- [6] E. Aboufadel, S. Schlicker (2003) "Wavelets : Wavelets for undergraduates." [Online]. Available: <http://www.gvsu.edu/math/wavelets/undergrad.htm>
- [7] P.V. O'Neil, *Advanced Engineering Mathematics*. 5<sup>th</sup> Ed. Thomson Brooks/Cole, 2003, pp. 841-854.
- [8] Chen, C.F. and Hsiao C.H., "Haar wavelet method for solving lumped and distributed-parameter systems." IEE Proc. Control Theory Appl. Vol 144, 1997, pp. 87-94
- [9] Chen, C.F. and Hsiao C.H., "Wavelet approach to optimizing dynamic systems." IEE Proc. Control Theory Appl. Vol 146, 1999, pp. 213-219.
- [10] Ülo Lepik, "Haar wavelet method for solving higher order differential equations" unpublished
- [11] Ülo Lepik, "Numerical solution of differential equations using Haar wavelets." Mathematics and Computers in Simulation vol 68, 2005, pp. 127-143.
- [12] Haydar Akca et.al., "Survey on wavelet transform and application in ODE and wavelet networks." unpublished
- [13] Hsiao C.H., "Haar wavelet approach to linear stiff systems." Mathematics and Computers in Simulation vol 64, 2004, pp. 561-567.